



Q) Why do we ALWAYS plot our data as the 1st Step?



Q) How can we be so sure about normality?



$$\bar{X} \rightarrow \mu$$



$$s^2 \rightarrow \sigma^2$$

$$\bar{X} = \mu \text{ normal}$$

$$\bar{X} \neq \mu \text{ skewed}$$

Report the Median


but what if you took 100 samples and plotted the means of those samples: sampling distribution



This idea is called the Central Limit Theorem:

if you take a sufficiently large # of random samples from a population, the distribution of the sample means will be approx normally distributed

Q) How do we interpret CLT?



if a sample, $\bar{X} \rightarrow \mu$

if a sampling distribution

$$E(\bar{X}) \rightarrow \mu \quad \mu_{\bar{X}} \rightarrow \mu$$

what about variance?

$$S^2 \rightarrow \sigma^2$$

$$S_{\bar{X}}^2 \rightarrow$$

variance?

$$VAR(\bar{X}) \rightarrow E(\sigma_{\bar{X}}^2) = \overset{\uparrow 2}{VAR}\left(\frac{X_1 + X_2 + \dots + X_n}{n}\right) \rightarrow \left(\frac{1}{n}\right)^2 \cdot VAR(\bar{X}_1 + \bar{X}_2 + \dots)$$

$$VAR(\bar{X}) = \frac{1}{n^2} \cdot (\sigma_1^2 + \sigma_2^2 + \dots + \sigma_n^2)$$

$$VAR(\bar{X}) = \frac{1}{n^2} \cdot \cancel{n} \sigma^2$$

$$VAR(\bar{X}) = \frac{\sigma^2}{n}$$

$$SD(\bar{X}) = \frac{\sigma}{\sqrt{n}} \rightarrow S_{\bar{X}} \rightarrow \sigma_{\bar{X}}$$


$\therefore$ SD of a sampling distribution is: standard error of the mean.

$$\sqrt{S^2} = S$$

why n ? $\uparrow n \quad \downarrow \text{SEM by } \sqrt{n}$

$$S > S_{\bar{X}}$$

Q) Why do we use normal? Probabilities



Total probability = 1

normal dist.

$$P(x = \#) = \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2}$$

$$\mu = 0, \sigma = 1$$

Standard normal distribution

if you do : Standard Normal Distribution $P(Z \leq \#)$

$$P(Z \leq 1.00) = .84$$

$$P(Z \geq 1.00) = 0.16$$

$$P(Z \leq -1.00) = 0.16$$

$$P(Z \geq -1.00) = 0.84$$

$$P(-1 \leq Z \leq 1) = 1 - 0.32 = 0.68$$

$$Z = P.0.05 (.025) \rightarrow P 0.975$$

$$P(Z \geq 1.96) = .05$$

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	.5000	.5040	.5080	.5120	.5160	.5199	.5239	.5279	.5319	.5359
0.1	.5398	.5438	.5478	.5517	.5557	.5596	.5636	.5675	.5714	.5753
0.2	.5793	.5832	.5871	.5910	.5948	.5987	.6026	.6064	.6103	.6141
0.3	.6179	.6217	.6255	.6293	.6331	.6368	.6406	.6443	.6480	.6517
0.4	.6554	.6591	.6628	.6664	.6700	.6736	.6772	.6808	.6844	.6879
0.5	.6915	.6950	.6985	.7019	.7054	.7088	.7123	.7157	.7190	.7224
0.6	.7257	.7291	.7324	.7357	.7389	.7422	.7454	.7486	.7517	.7549
0.7	.7580	.7611	.7642	.7673	.7704	.7734	.7764	.7794	.7823	.7852
0.8	.7881	.7910	.7939	.7967	.7995	.8023	.8051	.8078	.8106	.8133
0.9	.8159	.8186	.8212	.8238	.8264	.8289	.8315	.8340	.8365	.8389
1.0	.8413	.8438	.8461	.8485	.8508	.8531	.8554	.8577	.8599	.8621
1.1	.8643	.8665	.8686	.8708	.8729	.8749	.8770	.8790	.8810	.8830
1.2	.8849	.8869	.8888	.8907	.8925	.8944	.8962	.8980	.8997	.9015
1.3	.9032	.9049	.9066	.9082	.9099	.9115	.9131	.9147	.9162	.9177
1.4	.9192	.9207	.9222	.9236	.9251	.9265	.9279	.9292	.9306	.9319
1.5	.9332	.9345	.9357	.9370	.9382	.9394	.9406	.9418	.9429	.9441
1.6	.9452	.9463	.9474	.9484	.9495	.9505	.9515	.9525	.9535	.9545
1.7	.9554	.9564	.9573	.9582	.9591	.9599	.9608	.9616	.9625	.9633
1.8	.9641	.9649	.9656	.9664	.9671	.9678	.9686	.9693	.9699	.9706
1.9	.9713	.9719	.9726	.9732	.9738	.9744	.9750	.9756	.9761	.9767
2.0	.9772	.9778	.9783	.9788	.9793	.9798	.9803	.9808	.9812	.9817
2.1	.9821	.9826	.9830	.9834	.9838	.9842	.9846	.9850	.9854	.9857
2.2	.9861	.9864	.9868	.9871	.9875	.9878	.9881	.9884	.9887	.9890
2.3	.9893	.9896	.9898	.9901	.9904	.9906	.9909	.9911	.9913	.9916
2.4	.9918	.9920	.9922	.9925	.9927	.9929	.9931	.9932	.9934	.9936
2.5	.9938	.9940	.9941	.9943	.9945	.9946	.9948	.9949	.9951	.9952
2.6	.9953	.9955	.9956	.9957	.9959	.9960	.9961	.9962	.9963	.9964
2.7	.9965	.9966	.9967	.9968	.9969	.9970	.9971	.9972	.9973	.9974
2.8	.9974	.9975	.9976	.9977	.9977	.9978	.9979	.9979	.9980	.9981
2.9	.9981	.9982	.9982	.9983	.9984	.9984	.9985	.9985	.9986	.9986
3.0	.9987	.9987	.9987	.9988	.9988	.9989	.9989	.9989	.9990	.9990
3.1	.9990	.9991	.9991	.9991	.9992	.9992	.9992	.9992	.9993	.9993
3.2	.9993	.9993	.9994	.9994	.9994	.9994	.9994	.9995	.9995	.9995
3.3	.9995	.9995	.9995	.9996	.9996	.9996	.9996	.9996	.9996	.9997
3.4	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9998

Q) Application of CLT: confidence intervals (CI)

$$CI = \bar{X} \pm c \cdot \frac{\sigma}{\sqrt{n}}$$

$$= \bar{X} \pm z \cdot \frac{\sigma}{\sqrt{n}} \quad (\mu + \sigma \text{ known})$$

CI says: some % confident that the actual μ lies in that interval. μ

Ex: 68% CI ($\mu + \sigma$ known)

$$= \bar{X} \pm z \cdot \frac{\sigma}{\sqrt{n}}$$

what z corresponds to 68% probability?

$$\alpha = 1 - 0.68 = 0.32 \rightarrow \frac{\alpha}{2} = 0.16$$

$$= \bar{X} \pm 1 \cdot \frac{\sigma}{\sqrt{n}}$$

$$\therefore z \approx 0.84 : 1.00$$

$\therefore$ 68% CI is also the $S_{\bar{X}}$

Ex: 95% CI

$$= \bar{X} \pm z \cdot \frac{\sigma}{\sqrt{n}}$$

what value of z corresponds 95% probability

$$\alpha = 1 - 0.95 = .05 \rightarrow \frac{\alpha}{2} = 0.025$$

$$\bar{X} \pm 1.96 \cdot \frac{\sigma}{\sqrt{n}}$$


what $z \geq 0.975$? 1.96

in polling 95% CI $\approx$ margin of error ± 3

Q) Other distributions ?

TTable

Students "t"



a family of curves that correspond to a # of degrees of freedom (v)
 $d.o.f = n-1$

x, y, z
 $x+y+z = a$
 $x+y+z = b$
 $x+y+z = c$


Critical Values for Student's t Distribution

one-tail area	0.250	0.125	0.100	0.075	0.050	0.025	0.010	0.005	0.0005
two-tail area	0.500	0.250	0.200	0.150	0.100	0.050	0.020	0.010	0.0010
d.f. \ c	0.500	0.750	0.800	0.850	0.900	0.950	0.980	0.990	0.999
1	1.000	2.414	3.078	4.165	6.314	12.706	31.821	63.657	636.619
2	0.816	1.604	1.886	2.282	2.920	4.303	6.965	9.925	31.599
3	0.765	1.423	1.638	1.924	2.353	3.182	4.541	5.841	12.924
4	0.741	1.344	1.533	1.778	2.132	2.776	3.747	4.604	8.610
5	0.727	1.301	1.476	1.699	2.015	2.571	3.365	4.032	6.869
6	0.718	1.273	1.440	1.650	1.943	2.447	3.143	3.707	5.959
7	0.711	1.254	1.415	1.617	1.895	2.365	2.998	3.499	5.408
8	0.706	1.240	1.397	1.592	1.860	2.306	2.896	3.355	5.041
9	0.703	1.230	1.383	1.574	1.833	2.262	2.821	3.250	4.781
10	0.700	1.221	1.372	1.559	1.812	2.228	2.764	3.169	4.587
11	0.697	1.214	1.363	1.548	1.796	2.201	2.718	3.106	4.437
12	0.695	1.209	1.356	1.538	1.782	2.179	2.681	3.055	4.318
13	0.694	1.204	1.350	1.530	1.771	2.160	2.650	3.012	4.221
14	0.692	1.200	1.345	1.523	1.761	2.145	2.624	2.977	4.140
15	0.691	1.197	1.341	1.517	1.753	2.131	2.602	2.947	4.073
16	0.690	1.194	1.337	1.512	1.746	2.120	2.583	2.921	4.015
17	0.689	1.191	1.333	1.508	1.740	2.110	2.567	2.898	3.965
18	0.688	1.189	1.330	1.504	1.734	2.101	2.552	2.878	3.922
19	0.688	1.187	1.328	1.500	1.729	2.093	2.539	2.861	3.883
20	0.687	1.185	1.325	1.497	1.725	2.086	2.528	2.845	3.850
21	0.686	1.183	1.323	1.494	1.721	2.080	2.518	2.831	3.819
22	0.686	1.182	1.321	1.492	1.717	2.074	2.508	2.819	3.792
23	0.685	1.180	1.319	1.489	1.714	2.069	2.500	2.807	3.768
24	0.685	1.179	1.318	1.487	1.711	2.064	2.492	2.797	3.745
25	0.684	1.178	1.316	1.485	1.708	2.060	2.485	2.787	3.725
26	0.684	1.177	1.315	1.483	1.706	2.056	2.479	2.779	3.707
27	0.684	1.176	1.314	1.482	1.703	2.052	2.473	2.771	3.690
28	0.683	1.175	1.313	1.480	1.701	2.048	2.467	2.763	3.674
29	0.683	1.174	1.311	1.479	1.699	2.045	2.462	2.756	3.659
30	0.683	1.173	1.310	1.477	1.697	2.042	2.457	2.750	3.646
35	0.682	1.170	1.306	1.472	1.690	2.030	2.438	2.724	3.591
40	0.681	1.167	1.303	1.468	1.684	2.021	2.423	2.704	3.551
45	0.680	1.165	1.301	1.465	1.679	2.014	2.412	2.690	3.520
50	0.679	1.164	1.299	1.462	1.676	2.009	2.403	2.678	3.496
60	0.679	1.162	1.296	1.458	1.671	2.000	2.390	2.660	3.460
70	0.678	1.160	1.294	1.456	1.667	1.994	2.381	2.648	3.435
80	0.678	1.159	1.292	1.453	1.664	1.990	2.374	2.639	3.416
100	0.677	1.157	1.290	1.451	1.660	1.984	2.364	2.626	3.390
500	0.675	1.152	1.283	1.442	1.648	1.965	2.334	2.586	3.310
1000	0.675	1.151	1.282	1.441	1.646	1.962	2.330	2.581	3.300
$\infty (z_c)$	0.67	1.15	1.28	1.44	1.65	1.96	2.33	2.58	3.29


for degrees of freedom d.f. not in the table, use the closest d.f. that is smaller.

when $v = \infty$; $t \Rightarrow z$

F-Distribution (ANOVA, GLM, REGRESSION)



family of curves that correspond to a numerator & den. d.o.f



$n-1 : n=5$

df ₁	Numerator Degrees of Freedom (df ₁)															
df ₂	1	2	3	4	5	6	7	8	9	10	12	15	20	24	29	249.1
1	161.4	199.5	215.7	224.6	230.2	234.0	236.8	238.9	240.5	241.9	243.9	245.9	248.0	249.1		
2	18.51	19.00	19.16	19.25	19.30	19.33	19.35	19.37	19.38	19.40	19.41	19.43	19.45	19.45		
3	10.13	9.55	9.28	9.12	9.01	8.94	8.89	8.85	8.81	8.79	8.74	8.70	8.66	8.64		
4	7.71	6.94	6.59	6.39	6.26	6.16	6.09	6.04	6.00	5.96	5.91	5.86	5.80	5.77		
5	6.61	5.79	5.41	5.19	5.05	4.95	4.88	4.82	4.77	4.74	4.68	4.62	4.56	4.53		
6	5.99	5.14	4.76	4.53	4.39	4.28	4.21	4.15	4.10	4.06	4.00	3.94	3.87	3.84		
7	5.59	4.71	4.25	4.12	3.97	3.87	3.79	3.73	3.68	3.64	3.57	3.51	3.44	3.41		
8	5.32	4.46	4.07	3.84	3.69	3.58	3.50	3.44	3.39	3.35	3.28	3.22	3.15	3.12		
9	5.12	4.26	3.86	3.63	3.48	3.37	3.29	3.23	3.18	3.14	3.07	3.01	2.94	2.90		
10	4.96	4.10	3.71	3.48	3.33	3.22	3.14	3.07	3.02	2.98	2.91	2.85	2.77	2.74		
11	4.84	3.98	3.59	3.36	3.20	3.09	3.01	2.95	2.90	2.85	2.79	2.72	2.65	2.61		
12	4.75	3.89	3.49	3.26	3.11	3.00	2.91	2.85	2.80	2.75	2.69	2.62	2.54	2.51		
13	4.67	3.81	3.41	3.18	3.03	2.92	2.83	2.77	2.71	2.67	2.60	2.53	2.46	2.42		
14	4.60	3.74	3.34	3.11	2.96	2.85	2.76	2.70	2.65	2.60	2.53	2.46	2.39	2.35		
15	4.54	3.68	3.29	3.06	2.90	2.79	2.71	2.64	2.59	2.54	2.48	2.40	2.33	2.29		

Source: M. Meerrington and C. M. Thompson, "Tables of Percentage Points of the Inverted Beta (F) Distribution," *Biometrika*, vol. 33 (1943), pp. 73-88. Reproduced by permission of Oxford University Press and Biometrika trustees.

$n-1=7 : 8$